


Canine ACUTE PAIN SCALE



Pain Score	Example	Psychological & Behavioral	Response to Palpation	Body Tension
No Score		<input type="checkbox"/> Animal is sleeping and cannot be evaluated		
0		<input type="checkbox"/> Comfortable when resting <input type="checkbox"/> Happy, content <input type="checkbox"/> Not bothering wound or surgery site <input type="checkbox"/> Interested in or curious about surroundings	<input type="checkbox"/> Nontender to palpation of wound or surgery site, or to palpation elsewhere	Minimal
1		<input type="checkbox"/> Content to slightly unsettled or restless <input type="checkbox"/> Distracted easily by surroundings	<input type="checkbox"/> Reacts to palpation of wound, surgery site, or other body part by looking around, flinching, or whimpering	Mild
2		<input type="checkbox"/> Looks uncomfortable when resting. <input type="checkbox"/> May whimper or cry and may lick or rub wound or surgery site when unattended <input type="checkbox"/> Droopy ears, worried facial expression (arched eye brows, darting eyes) <input type="checkbox"/> Reluctant to respond when beckoned <input type="checkbox"/> Not eager to interact with people or surroundings but will look around to see what is going on	<input type="checkbox"/> Flinches, whimpers cries, or guards/pulls away	Mild to Moderate Reassess analgesic plan
3		<input type="checkbox"/> Unsettled, crying, groaning, biting or chewing wound when unattended <input type="checkbox"/> Guards or protects wound or surgery site by altering weight distribution (i.e., limping, shifting body position) <input type="checkbox"/> May be unwilling to move all or part of body	<input type="checkbox"/> May be subtle (shifting eyes or increased respiratory rate) if dog is too painful to move or is stoic <input type="checkbox"/> May be dramatic, such as a sharp cry, growl, bite or bite threat, and/or pulling away	Moderate Reassess analgesic plan
4		<input type="checkbox"/> Constantly groaning or screaming when unattended <input type="checkbox"/> May bite or chew at wound, but unlikely to move <input type="checkbox"/> Potentially unresponsive to surroundings <input type="checkbox"/> Difficult to distract from pain	<input type="checkbox"/> Cries at non-painful palpation (may be experiencing allodynia, wind-up, or fearful that pain could be made worse) <input type="checkbox"/> May react aggressively to palpation	Moderate to Severe May be rigid to avoid painful movement Reassess analgesic plan



Comments _____

Dr. Michelle V. Rose DVM, CCRP
 (415) 326-4838 • mrose@mrosevet.com
 www.mrosevet.com